



## Product Datasheet

### HPRT antibody GRP140

<b>Description</b>	HPRT1 has a central role in the generation of purine nucleotides through the purine salvage pathway. HPRT1 catalyzes conversion of hypoxanthine to inosine monophosphate and guanine to guanosine monophosphate via transfer of the 5-phosphoribosyl group from 5-phosphoribosyl 1-pyrophosphate (Keebaugh et al., 2007 [PubMed 16928426]).[supplied by OMIM]
<b>Species/Host</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat
<b>Conjugation</b>	Unconjugated
<b>Tested Applications</b>	ICC, IF, IHC-P, WB
<b>Immunogen</b>	Recombinant protein encompassing a sequence within the center region of human HPRT. The exact sequence is proprietary.
<b>Form/Appearance</b>	Liquid: 1XPBS, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative.
<b>Concentration</b>	0.65 mg/ml
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Note</b>	For research use only.
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Purified by antigen-affinity chromatography.
<b>Uniprot ID</b>	<b>P00492</b>
<b>Entrez</b>	<b>3251</b>
<b>Dilution Range</b>	WB: 1:500-1:3000, ICC: 1:100-1:1000, IHC-P: 1:100-1:1000



Various whole cell extracts (30 µg) were separated by 12% SDS-PAGE, and the membrane was blotted with HPRT antibody (GRP592) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.



HPRT antibody detects HPRT protein at cytoplasm in rat colon by immunohistochemical analysis. Sample: Paraffin-embedded rat colon. HPRT antibody (GRP592) diluted at 1:500.