

## Product Datasheet

### Mouse IgG2a Isotype Control Purified Low Endotoxin GRP11908

|                            |   |
|----------------------------|---|
| <b>Description</b>         | This mouse IgG2a monoclonal antibody (clone MOPC-173) reacts with an unknown epitope. It does not react with a variety of resting, activated, live, and fixed mouse, rat and human tissues.   |
| <b>Tested Applications</b> | ELISA, FC, FA, ICC, IHC-Fr, IHC-P, IP, WB   |
| <b>Immunogen</b>           | The transplantable plasmacytoma MOPC-173 was induced by intraperitoneal injection of mineral oils into BALB/c mice.   |
| <b>Concentration</b>       | 1 mg/ml   |
| <b>Storage</b>             | Store at 2-8°C. Do not freeze. Do not use after expiration date stamped on the label.   |
| <b>Note</b>                | For research use only.  |
| <b>Isotype</b>             | Mouse IgG2a kappa   |
| <b>Clonality</b>           | Monoclonal  |
| <b>Clone ID</b>            | MOPC-173  |
| <b>Application Notes</b>   | Negative control: The reagent is intended as an isotype control to establish the amount of non-specific antibody binding. For your particular experiment, use the same concentration of this control antibody as the recommended working concentration of the antigen-specific antibody. Also, when working with prediluted antibodies, dilute the isotype control to the same concentration as is the concentration of the antigen-specific antibody in the prediluted antibody solution you are using. If under particular experimental conditions the background signal of the isotype control is too high (usually when working concentrations of used antibodies are above 10 µg/ml of incubation mixture), change the conditions of your experiment to reduce the background. |