

Product Datasheet

Rabbit anti-Goat IgG Fc, HRP conjugated GRP12266

Species/Host	Rabbit
Form/Appearance	Lyophilized
Storage	Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified rabbit IgG
Dilution Range	1 : 500-1 : 5000 (IHC), 1 : 10 000-1 : 50 000 (WB)
Application Notes	Additional Information: This antibody reacts with the heavy chains on goat IgG based on immunoelectrophoresis. Minimum cross-reactivity is observed to the light chains or to non-immunoglobulin goat serum proteins based on immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free and 0.1 % (v/v) Kathon CG is used as preservative. Use of sodium azide will inhibit enzymatic activity of horseradish peroxidase. BSA and milk have to be replaced by other blocking reagents, like donkey serum or commercial formulations which are free from bovine IgG. Background: Rabbit anti-goat IgG Fc is a secondary antibody to goat Fc part conjugated to HRP. Antibodies are affinity purified using solid phase goat IgG. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.