

German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

Donkey anti-Rabbit IgG (H&L), HRP conjugated, min. cross-reactivity to bovine, chicken, goat, guinea pig, hamster, horse, human, mouse, rat, sheep IgG **GRP12270**

Species/Host Donkey

Form/Appearance Lyophilized

Store lyophilized material at 2-8°C. For long time storage after **Storage**

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

Note For research use only.

Clonality Polyclonal

Purity Affinity purified IgG

The optimal working dilution should be determined by the **Dilution Range**

investigator.

Application Notes Additional Information: This antibody reacts with the heavy chains on rabbit IgG

and with the light chains on all rabbit immunoglobulins based on immunoelectrophoresis. Minimum cross-reactivity is observed to

non-immunoglobulin rabbit serum proteins based on

immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free and 0.1 % (v/v) Kathon CG is used as preservative. Use of sodium azide will inhibit enzymatic activity of horseradish peroxidase. Background: Donkey anti-rabbit $\lg G$ (H&L) is a secondary antibody, HRP conjugated, which binds to all rabbit $\lg Gs$ in

immunological assays. Antibody is specially adsorbed against

bovine, chicken, goat, guinea pig, hamster, horse, human, mouse, rat, sheep IgG. Antibodies are affinity purified using solid phase rabbit IgG. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room

temperature to dissolve. Prepare fresh working dilutions daily.

German Research Products - GRP GmbH D-85410 Haag/Amper, Germany Email: info@grp-ak.de | Phone: +49 (0)8167 6335