

## **Product Datasheet**

## Goat anti-Rabbit IgG (H&L), F(ab)'2 fragment, FITC conjugated GRP12272

Species/Host	Goat
Form/Appearance	Liquid
Storage	Store at 2-8°C. Shelf life of this product is one year from date of receipt.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified IgG, F(ab)'2 fragment
Dilution Range	1:50-1:5000 (ICC) (Flow cyt), 1:20-1:2000 (IHC)
Application Notes	Additional Information: This antibody reacts with the heavy chains on rabbit IgG and with the light chains on all rabbit immunoglobulins based on immunoelectrophoresis.Minimum cross-reactivity is observed to non-immunoglobulin rabbit serum proteins based on immunoelectrophoresis.Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative. Background: Goat anti-rabbit IgG (H&L), F(ab)'2 fragment of FITC (fluorescein-5-isothiocyanate) conjugated secondary antibody which binds to rabbit IgG in immunological assays. FITC has Amax = 494 nm, Emax = 518 nm. Fluor to protein ratio is 3-7 moles FITC per mole antibody. Antibody purity is >90% based on SDS-PAGE and is affinity purified on solid phase rabbit IgG. Antibody solution may contain small amounts of intact IgG.