

Product Datasheet

Goat anti-Human IgM (μ chain), F(ab)'2 fragment, TRITC conjugated, min. cross-reactivity to human IgG or IgA GRP12330

Species/Host	Goat
Form/Appearance	Liquid
Storage	Store at 2-8°C. Shelf life of this product is one year from date of receipt.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified IgG
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: This antibody reacts with the mu chains on human IgM based on immunoelectrophoresis. Minimum cross-reactivity is observed with IgG, F(ab)'2 fragment, IgG or IgA based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin human serum proteins based on immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative. Background: Goat anti-human IgM (μ chain), Rhodamine (TRITC-tetramethylrhodamine-5-isothiocyanate) conjugated, is a F(ab)'2 fragment of a secondary antibody which binds human IgM in immunological assays. TRITC has λ_{max} = 550 nm, λ_{em} = 570 nm. Antibody is adsorbed against human IgG or IgA and affinity purified using solid phase human IgM.</p>