

## Product Datasheet

### Goat anti-Rat IgG (H&L), HRP conjugated, min. cross-reactivity to human, mouse IgG GRP12334

<b>Species/Host</b>	Goat
<b>Form/Appearance</b>	Lyophilized
<b>Storage</b>	Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
<b>Note</b>	For research use only.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified IgG
<b>Dilution Range</b>	The optimal working dilution should be determined by the investigator.
<b>Application Notes</b>	<p>Additional Information: This antibody reacts with the heavy chains on rat IgG and with the light chains on all rat immunoglobulins based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin rat serum proteins or IgG from human or mouse. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free and 0.1 % (v/v) Kathon CG is used as preservative.</p> <p>Background: Goat anti-rat IgG (H&amp;L), HRP conjugated, is a secondary antibody which binds to rat IgGs in immunological assays. Antibodies are adsorbed against human, mouse IgG and affinity purified using solid phase rat IgG.</p> <p>Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.</p>