

Product Datasheet

Rabbit anti-Goat IgG Fc, TRITC conjugated GRP12341

Species/Host	Rabbit
Form/Appearance	Liquid
Storage	Store at 2-8°C. Shelf life of this product is one year from date of receipt.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified IgG
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: This antibody reacts with the heavy chains on goat IgG based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin goat serum proteins or immunoglobulin light chains based on immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative. BSA and milk have to be replaced by other blocking reagents, like donkey serum or commercial formulations which are free from bovine IgG. Background: Rabbit anti-goat IgG Fc, Rhodamine (TRITC-tetramethylrhodamine-5-isothiocyanate) conjugated, is a secondary antibody which binds to goat IgG Fc constant part in immunological assays. TRITC has Amax = 550 nm, Emax = 570 nm. Antibodies are affinity purified on solid phase goat IgG Fc.</p>