

Product Datasheet

Chicken anti-Goat IgG (H&L), TRITC conjugated GRP12370

Species/Host	Chicken
Form/Appearance	Liquid
Storage	Store at 2-8°C. Shelf life of this product is one year from date of receipt.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified IgY
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: This antibody reacts with the heavy chains on goat IgG and with the light chains on all goat immunoglobulins based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin goat serum proteins based on immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative. BSA and milk have to be replaced by other blocking reagents, like donkey serum or commercial formulations which are free from bovine IgG. Background: Chicken anti-goat IgG (H&L), Rhodamine (TRITC-tetramethylrhodamine-5-isothiocyanate) conjugated is a secondary antibody which binds to goat IgGs in immunological assays. TRITC has Amax = 550 nm, Emax = 570 nm. Antibodies are affinity purified using solid phase goat IgG.</p>