

## Product Datasheet

### Donkey anti-Rabbit IgG (H&L), TRITC conjugated, min. cross-reactivity to mouse IgG/serum GRP12377

|                          |  |
|--------------------------|--|
| <b>Species/Host</b>      | Donkey   |
| <b>Form/Appearance</b>   | Liquid   |
| <b>Storage</b>           | Store at 2-8°C. Shelf life of this product is one year from date of receipt.   |
| <b>Note</b>              | For research use only.   |
| <b>Clonality</b>         | Polyclonal   |
| <b>Purity</b>            | Affinity purified IgG  |
| <b>Dilution Range</b>    | The optimal working dilution should be determined by the investigator.   |
| <b>Application Notes</b> | <p>Additional Information: This antibody reacts with the heavy chains on rabbit IgG and with the light chains on all rabbit immunoglobulins based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin rabbit serum proteins, mouse IgG or mouse serum based on immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative. Background: Donkey anti-rabbit IgG (H&amp;L), Rhodamine (TRITC-tetramethylrhodamine-5-isothiocyanate) conjugated, is a secondary antibody which binds to rabbit IgGs in immunological assays. TRITC has <math>\lambda_{max} = 550</math> nm, <math>\lambda_{em} = 570</math> nm. Antibodies are adsorbed against mouse IgG/serum and affinity purified using solid phase rabbit IgG.</p> |