

## Product Datasheet

### Goat anti-Human IgA + IgG + IgM, F(ab)'2 fragment, Affinity purified, Unconjugated GRP12407

|                          |  |
|--------------------------|--|
| <b>Species/Host</b>      | Goat   |
| <b>Form/Appearance</b>   | Liquid   |
| <b>Storage</b>           | Store at 2-8°C.  |
| <b>Note</b>              | For research use only.   |
| <b>Clonality</b>         | Polyclonal   |
| <b>Purity</b>            | Affinity purified IgG, F(ab)'2 fragment  |
| <b>Dilution Range</b>    | The optimal working dilution should be determined by the investigator.   |
| <b>Application Notes</b> | <p>Additional Information: This antibody reacts with the heavy and light chains on human IgA + IgG + IgM based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin human serum proteins based on immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, H 7, 0.05 % (w/v) sodium azide as preservative.</p> <p>Background: Goat anti-human IgA + IgG + IgM, affinity purified F(ab)'2 fragment, is a secondary antibody which binds to human IgA and IgG and IgM in immunological assays. Antibody purity is &gt; 90% based on SDS-PAGE. Antibody solution may contain small amounts of intact IgG.</p> |