

## Product Datasheet

### Rabbit anti-Goat IgG Fc, Affinity purified, Unconjugated, min. cross-reactivity to human serum GRP12472

<b>Species/Host</b>	Rabbit
<b>Form/Appearance</b>	Liquid
<b>Storage</b>	Store at 2-8°C.
<b>Note</b>	For research use only.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified IgG
<b>Dilution Range</b>	The optimal working dilution should be determined by the investigator.

**Application Notes** Additional Information: This antibody reacts with the heavy chains on goat IgG based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin goat serum proteins or immunoglobulin light chains based on immunoelectrophoresis. Minimum cross-reactivity is observed with human serum proteins based on immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7, 0.05 % (w/v) sodium azide as preservative. BSA and milk have to be replaced by other blocking reagents, like donkey serum or commercial formulations which are free from bovine IgG. Background: Rabbit anti-goat IgG Fc, affinity purified, is a secondary antibody which binds to goat IgG Fc constant part in immunological assays. Antibodies are adsorbed against human serum and are affinity purified using solid phase goat IgG.