

Product Datasheet

Goat anti-Human IgM (μ chain), TRITC conjugated GRP12485

Species/Host	Goat
Form/Appearance	Liquid
Storage	Store at 2-8°C. Shelf life of this product is one year from date of receipt.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified IgG
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: This antibody reacts with the mu chains on human IgM based on immunoelectrophoresis. Minimum cross-reactivity is observed with IgG, F(ab)'2 fragment, IgA or IgG based on immunoelectrophoresis. Minimum cross-reactivity is observed to non-immunoglobulin human serum proteins based on immunoelectrophoresis. Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative. Background: Goat anti-human IgM (μ chain), Rhodamine (TRITC -tetramethylrhodamine-5-isothiocyanate) conjugated is a secondary antibody which binds to human IgM in immunological assays. TRITC Amax = 550 nm, Emax = 570 nm. Antibodies affinity purified using solid phase human IgM.</p>