

## Product Datasheet

### Goat anti-Human IgA + IgG + IgM, F(ab)'2 fragment, TRITC conjugated GRP12496

|                          |   |
|--------------------------|---|
| <b>Species/Host</b>      | Goat  |
| <b>Form/Appearance</b>   | Liquid  |
| <b>Storage</b>           | Store at 2-8°C. Shelf life of this product is one year from date of receipt.  |
| <b>Note</b>              | For research use only.  |
| <b>Clonality</b>         | Polyclonal  |
| <b>Purity</b>            | Affinity purified IgG, F(ab)'2 fragment   |
| <b>Dilution Range</b>    | The optimal working dilution should be determined by the investigator.  |
| <b>Application Notes</b> | <p>Additional Information: This antibody reacts with the heavy and light chains on human IgG, IgA and IgM based on immunoelectrophoresis. No reactivity is observed to non-immunoglobulin human serum proteins based on immunoelectrophoresis. Background: Goat anti-human IgA + IgG + IgM, Rhodamine (TRITC-tetramethylrhodamine-5-isothiocyanate) conjugated is a F(ab)'2 fragment of a secondary antibody which binds to human IgA and IgG and IgM in immunological assays. TRITC has <math>\lambda_{max} = 550 \text{ nm}</math>, <math>\lambda_{em} = 570 \text{ nm}</math>. Antibody purity is &gt; 90% based on SDS-PAGE. Antibody solution may contain small amounts of intact IgG.</p> |