

Product Datasheet

S100 beta antibody GRP158

Description

The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21; however, this gene is located at 21q22.3. This protein may function in Neurite extension, proliferation of melanoma cells, stimulation of Ca²⁺ fluxes, inhibition of PKC-mediated phosphorylation, astrocytosis and axonal proliferation, and inhibition of microtubule assembly. Chromosomal rearrangements and altered expression of this gene have been implicated in several neurological, neoplastic, and other types of diseases, including Alzheimer's disease, Down's syndrome, epilepsy, amyotrophic lateral sclerosis, melanoma, and type I diabetes. [provided by RefSeq, Jul 2008]

Species/Host

Rabbit

Reactivity

Human, Mouse, Rat, Squirrel

Conjugation

Unconjugated

Tested Applications

ICC, IF, IHC-Fr, IHC-P, WB

Immunogen

Full length human S100 beta Recombinant protein.

Form/Appearance

Liquid: 1XPBS, 20% Glycerol (pH7). 0.025% ProClin 300 was added as a preservative.

Concentration

1.05 mg/ml

Storage

Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

Note

For research use only.

Isotype

IgG

Clonality

Polyclonal

Purity

Purified by antigen-affinity chromatography.

Uniprot ID


P04271

Entrez


6285

Dilution Range

WB: 1:500-1:3000, IHC-P: 1:100-1:1000, IHC-Fr: 1:100-1:1000



S100B antibody detects S100B protein at glial cell on mouse hind brain by immunohistochemical analysis.
Sample: Paraffin-embedded mouse hind brain. S100B antibody (GRP610) dilution: 1:500.



Whole cell extract (30 µg) was separated by 15% SDS-PAGE, and the membrane was blotted with S100B antibody (GRP610) diluted at 1:500.