

German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

Sheep anti-Rabbit IgG (H&L), HRP conjugated GRP12577

Species/Host Sheep

Immunogen Purified rabbit IgG (H&L)

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C. For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

Note For research use only.

Clonality Polyclonal

Purity Affinity purified sheep IgG

Dilution Range This conjugate is suitable for all immunoassay applications. The

optimal working dilution should be determined by the investigator. Suggested starting dilution(s): 1:500-1:5000

(IHC), 1 : 200-1 : 5000 (WB)

Application Notes Additional Information: Purity of this preparation is > 95% based on SDS-PAGE.

Antibody concentration is 1.0 mg/ml. Antibody is supplied in 10 mM sodium phosphate, 0.15 M sodium chloride, pH 7.2.1 % (w/v) B, Protease/lgG free. Contains 0.1 % (v/v) Kathon CG as preservative of bacterial growth.Based on immunoelectrophoresis, this antibody reacts with: heavy chains on rabbit lgG, light chains on all rabbit immunoglobulins. Based on immunoelectrophoresis, no

reactivity is observed to: non-immunoglobulin rabbit serum proteins.

Background: Sheep anti-rabbit IgG (H&L) is a secondary antibody conjugated to HRP, which binds to rabbit IgG (H&L) in immunological assays. Antibody is affinity purified using solid phase rabbit IgG (H&L). Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room

temperature to dissolve. Prepare fresh working dilutions daily.