German Research Products - GRP GmbH
In der Stockwiese 26
D-85410 Haag/Amper, Germany
Email: info@grp-ak.de
Phone: +49 (0)8167 6335

Product Datasheet

Goat anti-Human IgA heavy (alpha chain), F(ab)'2 fragment, HRP conjugated

GRP12631

Species/Host	Goat
Reactivity	Human
Predicted Reactivity	Human IgA Heavy chain (alpha) F(ab)'2 fragment
Tested Applications	ELISA, WB, IHC
Immunogen	purified human IgA, heavy (alpha chain)
Form/Appearance	Lyophilized
Storage	Store lyophilized material at 2-8°C.For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 0.5 mg of antibody in 0.55 ml of sterile water add 0.55 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified goat IgG
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	Additional Information: No reactivity is observed to the light chains or non-immunoglobulin human serum proteins. HRP-conjugate is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free0.1 % (v/v) of Kathon CG is used as preservative. Use of sodium azide will inhibit enzyme activity of horseradish peroxidasePurity is ≥ 90% based on SDS-PAGE. May contain small amounts of intact IgG. Background: Goat anti-human IgA heavy (alpha chain) is a secondary antibody conjugated to HRP which binds to human IgA heavy (alpha chain) F(ab)'2 fragments in immunological assays. Reconstitution: For reconstitution add 0.55 ml of sterile water