

Product Datasheet

Goat anti-Human IgG Fc, Biotin conjugated, min. cross-reactivity to human IgA+IgM GRP12670

| | |
|-----------------------------|---|
| Species/Host | Goat |
| Reactivity | Human |
| Predicted Reactivity | Human IgG Fc (two Heavy chains with constant domains) |
| Immunogen | Purified human IgG |
| Form/Appearance | Liquid |
| Storage | Store non-diluted antibody at 2-8°C. Shelf life is 1 year from date of receipt. |
| Note | For research use only. |
| Clonality | Polyclonal |
| Purity | Affinity purified goat IgG |
| Dilution Range | The optimal working dilution should be determined by the investigator. |
| Application Notes | <p>Additional Information: This antibody reacts with the heavy chains on human IgG based on immunoelectrophoresis. No reactivity is observed to immunoglobulin light chains, and human IgA and IgM based on immunoelectrophoresis. Biotin conjugate is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. Contains 0.05 % (w/v) sodium azide as preservative. Background: Goat anti-human IgG Fc (two heavy chains with constant domains) is a secondary antibody conjugated to biotin which binds to human IgG Fc (two heavy chains with constant domains) in immunological assays.</p> |