

Product Datasheet

Goat anti-Human IgG Fc, Biotin conjugated, min. cross-reactivity to human IgA+IgM GRP12670

Species/Host	Goat
Reactivity	Human
Predicted Reactivity	Human IgG Fc (two Heavy chains with constant domains)
Immunogen	Purified human IgG
Form/Appearance	Liquid
Storage	Store non-diluted antibody at 2-8°C. Shelf life is 1 year from date of receipt.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified goat IgG
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: This antibody reacts with the heavy chains on human IgG based on immunoelectrophoresis. No reactivity is observed to immunoglobulin light chains, and human IgA and IgM based on immunoelectrophoresis. Biotin conjugate is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. Contains 0.05 % (w/v) sodium azide as preservative Background: Goat anti-human IgG Fc (two heavy chains with constant domains) is a secondary antibody conjugated to biotin which binds to human IgG Fc (two heavy chains with constant domains) in immunological assays.</p>