

Product Datasheet

Goat anti-Human IgG Fc, Biotin conjugated, F(ab)'2 fragment GRP12673

Species/Host	Goat
Reactivity	Human
Predicted Reactivity	Human IgG Fc
Immunogen	purified human IgG (H&L)
Form/Appearance	Liquid
Storage	Store non-diluted antibody at 2-8°C. Shelf life is 1 year from date of receipt.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified goat IgG
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: This antibody reacts with the heavy chains on human IgG based on immunoelectrophoresis. No reactivity is observed with the light chains on human immunoglobulins and non-immunoglobulin human serum proteins and human IgG F(ab)'2 fragment based in immunoelectrophoresis. Biotin conjugate is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. Contains 0.05 % (w/v) sodium azide as preservative. Purity is ≥ 90% based on SDS-PAGE. May contain small amounts of intact IgG. Background: Goat anti-human IgG Fc, F(ab)'2 fragment is a secondary antibody conjugated to biotin which binds to human IgG Fc fragment in immunological assays.</p>