

## Product Datasheet

### Goat anti-Human IgG Fc, F(ab)'2 fragment, Affinity purified, Unconjugated, min. cross-reactivity to bovine/mouse/rabbit serum GRP12675

Species/Host	Goat
Reactivity	Human
Predicted Reactivity	Human IgG Fc (two Heavy chains with constant domains)
Tested Applications	ELISA
Immunogen	Purified human IgG
Form/Appearance	Liquid
Storage	Store at 2-8°C. Shelf life is one year from date of receipt.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified goat IgG
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: This antibody reacts with the heavy chains on human IgG based on immunoelectrophoresis. No reactivity is observed with the light chains on human immunoglobulins or non-immunoglobulin human serum or bovine, mouse or rabbit serum proteins or human F(ab)'2 fragment based on immunoelectrophoresis. This antibody has been absorbed against solid phase normal bovine, mouse and rabbit serum in addition to mouse and rabbit IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 0.05 % (w/v) of Sodium azide as preservative. Antibody concentration is 3 mg/ml. Background: Goat anti-human IgG Fc (two heavy chains with constant domains) F(ab)'2 fragment is a secondary antibody which binds to human IgG Fc (two heavy chains with constant domains) in immunological assays.</p>