

German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

Chicken anti-Rabbit IgG (H&L), HRP conjugated, min. cross-reactivity to human, mouse IgG GRP12692

Species/Host Chicken

Reactivity Rabbit

Predicted Reactivity Rabbit IgG (H&L)

Tested Applications ELISA, WB, IHC

Immunogen Purified Rabbit IgG

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C.For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 0.5 mg of antibody in 0.55 ml of sterile water add 0.55 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to

mix well but without foaming.

Note For research use only.

Clonality Polyclonal

Purity Affinity purified chicken IgY

Dilution Range The optimal working dilution should be determined by the

investigator.

Application Notes Additional Information: This antibody reacts with the heavy chains on rabbit IgG

and with the light chains on all rabbit immunoglobulins based on

immunoelectrophoresis.No reactivity is observed to non-immunoglobulin rabbit

serum proteins and human and mouse serum and IgG on

immunoelectrophoresis. This antibody was absorbed against solid phase human and mouse serum and IgG. HRP-conjugate is supplied in 10 mM Sodium

and mouse serum and IgG. HRP-conjugate is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free0.1 % (v/v) of Kathon CG is used as preservative. Use of sodium azide will inhibit enzyme activity of horseradish peroxidase Background: Chicken

inhibit enzyme activity of horseradish peroxidase Background: Chicken

anti-rabbit IgG (H&L) is a secondary antibody conjugated to HRP which binds to rabbit IgG (H&L) in immunological assays. Reconstitution: For reconstitution add 0.55 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve.

Prepare fresh working dilutions daily.