

**German Research Products - GRP GmbH** 

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

## **Product Datasheet**

## Goat anti-Rabbit IgG Fc, HRP conjugated GRP12704

Species/Host Goat

Reactivity Rabbit

Predicted Reactivity Rabbit IgG Fc (two Heavy chains with constant domains)

Tested Applications ELISA, WB, IHC

Immunogen Purified Rabbit IgG

Form/Appearance Lyophilized

**Storage** Store lyophilized material at 2-8°C.For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

**Note** For research use only.

**Clonality** Polyclonal

**Purity** Affinity purified goat IgG

**Dilution Range** The optimal working dilution should be determined by the

investigator.

**Application Notes** Additional Information: This antibody reacts with the heavy chains on rabbit IgG

based on immunoelectrophoresis. No reactivity is observed to the light chains on rabbit immunoglobulins or non-immunoglobulin rabbit serum proteins based on immunoelectrophoresis. This antibody will cross react with the heavy chain on other species IgG, but not with IgM or IgA. HRP-conjugate is supplied in 10 mM

Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 % (w/v) BSA,

Protease/IgG free0.1 % (v/v) of Kathon CG is used as preservative. Use of sodium azide will inhibit enzyme activity of horseradish peroxidase Background: Goat anti-rabbit IgG Fc (two heavy chains with constant domains) is a secondary antibody conjugated to HRP which binds to rabbit IgG Fc (two heavy chains with constant domains) in immunological assays. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to

dissolve. Prepare fresh working dilutions daily.