

German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

Rabbit anti-Goat IgG (H&L), TRITC conjugated GRP12732

Species/Host Rabbit

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C.For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

Note For research use only.

Clonality Polyclonal

Purity Affinity purified rabbit IgG

Dilution Range The optimal working dilution should be determined by the

investigator. (ICC), immunohistochemistry (IHC) (Flow cyt)

Application Notes Additional Information: This antibody reacts with the heavy chains on goat IgG

and with the light chains on all goat immunoglobulins based on immunoelectrophoresis. Minimum cross-reactivity is observed to

non-immunoglobulin goat serum proteins based on

immunoelectrophoresis.Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.BSA and milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free from bovine IgG. Background: Rabbit anti-goat IgG (H&L), is a Rhodamine (TRITC) conjugated secondary antibody which binds to goat IgG (H&L) in immunological assays. Rhodamine - Tetramethylrhodamine-5-isothiocyanate

immunological assays. Rhodamine - Tetramethylrhodamine-5-isothiocyanate (TRITC) has Amax = 550 nm, Emax = 570 nm. Antibodies are affinity purified using solid phase goat IgG. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare

fresh working dilutions daily.