In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

beta Tubulin 3/ Tuj1 antibody **GRP160**

This gene encodes a class III member of the beta tubulin protein Description

family. Beta tubulins are one of two core protein families (alpha and beta tubulins) that heterodimerize and assemble to form microtubules. This protein is primarily expressed in neurons and may be involved in neurogenesis and axon guidance and maintenance. Mutations in this gene are the cause of congenital fibrosis of the extraocular muscles type 3. Alternate splicing results in multiple transcript variants. A pseudogene of this gene

is found on chromosome 6. [provided by RefSeq]

Species/Host Rabbit

Reactivity Human, Mouse, Rat

Conjugation Unconjugated

Tested Applications ICC, IF, IHC-Fr, IHC-P, IP, WB

Carrier-protein conjugated synthetic peptide encompassing a **Immunogen**

sequence within the center region of human beta Tubulin 3/ TUJ1

The exact sequence is proprietary.

Liquid: 1XPBS, 20% Glycerol (pH7). 0.025% ProClin 300 was Form/Appearance

added as a preservative.

Concentration 0.1 mg/ml

Storage Store as concentrated solution. Centrifuge briefly prior to opening

vial. For short-term storage (1-2 weeks), store at 4°C. For

long-term storage, aliquot and store at -20°C or below. Avoid

multiple freeze-thaw cycles.

For research use only. Note

Isotype **IgG**

Clonality Polyclonal


Purified by antigen-affinity chromatography. **Purity**

Uniprot ID Q13509

Entrez 10381

Dilution Range WB: 1:500-1:50000,ICC: 1:100-1:1000,IHC-P:

1:100-1:1000,IHC-Fr: 1:100-1:1000,IP: 1:100-1:500



beta Tubulin 3/TUJ1 antibody detects beta Tubulin 3/

TUJ1

protein by western blot analysis. Various whole

cell


extracts (30 ?g) were separated by 10% SDS-PAGE, and

the membrane was blotted with

beta Tubulin 3/ TUJ1 antibody

(GRP612) diluted at a

dilution



beta Tubulin 3/TUJ1 antibody detects beta Tubulin 3/ TUJ1

protein at cytoplasm in rat brain by immunohistochemical analysis. Sample: Paraffin-embedded rat brain.

beta Tubulin 3/ TUJ1 antibody

(GRP612) diluted at 1:500.