

**German Research Products - GRP GmbH** 

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

## **Product Datasheet**

## Rabbit anti-Hamster IgG (H&L), HRP conjugated GRP12770

Species/Host Rabbit

Immunogen Purified hamster IgG (H&L)

Form/Appearance Lyophilized

**Storage** Store lyophilized material at 2-8°C.For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

**Note** For research use only.

**Clonality** Polyclonal

Purity Affinity purified rabbit IgG

**Dilution Range** This conjugate is suitable for all immunoassay applications. The

optimal working dilution should be determined by the

investigator.

**Application Notes** Additional Information: Purity of this preparation is > 95% based on SDS-PAGE.

Antibody concentration is 1.0 mg/ml. Antibody is supplied in 10 mM sodium phosphate, 0.15 M sodium chloride, pH 7.2.1 % (w/v) B, Protease/IgG free. Contains 0.1 % (v/v) Kathon CG as preservative of bacterial growth.Based on immunoelectrophoresis, this antibody reacts with: heavy chains on hamster IgG, light chains on all hamster immunoglobulins. Based on immunoelectrophoresis, no reactivity is observed to: non-immunoglobulin hamster serum proteins Background: Rabbit anti-hamster IgG (H&L) is a secondary antibody conjugated to HRP, which binds tohamster IgG (H&L) in immunological assays. Antibody is affinity purified using solid phase hamster IgG (H&L). Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room

temperature to dissolve. Prepare fresh working dilutions daily.