

German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

Goat anti-Mouse IgG (H&L), HRP conjugated GRP12771

Species/Host Goat

Immunogen Purified mouse IgG (H&L) AAA51107

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C.For long time storage after

reconstitution, dilute the antibody solution with glycerol to a fina concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

Note For research use only.

Clonality Polyclonal

Purity Affinity purified IgG

Dilution Range This conjugate is suitable for all immunoassay applications. The

optimal working dilution should be determined by the

investigator.

Application Notes Additional Information: Purity of this preparation is > 95% based on SDS-PAGE.

Antibody concentration is 1.0 mg/ml. Antibody is supplied in 10 mM sodium phosphate, 0.15 M sodium chloride, pH 7.2.1 % (w/v) B, Protease/IgG free. Contains 0.1 % (v/v) Kathon CG as preservative of bacterial growth.Based on immunoelectrophoresis, this antibody reacts with:heavy chains on mouse IgG, light chains on all mouse immunoglobulins. Based on immunoelectrophoresis, no reactivity is observed to: non-immunoglobulin mouse serum proteins.The antibody will detect all isotypes of mouse IgG. Background: Goat anti-mouse IgG (H&L) is a secondary antibody conjugated to HRP, which binds to mouse IgG

(H&L) in immunological assays. Antibody is affinity purified using solid phaseMouse IgG (H&L). Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh

working dilutions daily.

