German Research Products - GRP GmbH
In der Stockwiese 26
D-85410 Haag/Amper, Germany
Email: info@grp-ak.de
Phone: +49 (0)8167 6335

Product Datasheet

Goat anti-Chicken IgY (H&L), DyLight® 633 conjugated
GRP12782

Species/Host	Goat
Reactivity	Chicken
Predicted Reactivity	Chicken IgY Heavy and Light chains (H&L)
Tested Applications	ICC, IHC
Immunogen	purified chicken IgY, whole molecule
Form/Appearance	Lyophilized
Storage	Store lyophilized material at 2-8°C. Product is stable for 4 weeks at 2-8°C after rehydration.For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified goat IgG
Dilution Range	1 : 50 -1 : 5 000 (ICC), 1 : 20- 1 : 2000 (IHC)
Application Notes	Additional Information: Based on immunoelectrophoresis this antibody reacts with heavy chains on chicken IgG and light chains on all chicken immunoglobulins.No reactivity is observed to non-immunoglobulin chicken serum proteins based in immunoelectrophoresis. Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative.DyLight® 633 (Ex = 638 nm; Em = 658 nm). Background: Goat anti-chicken IgY, DyLight® 633 Conjugate is a secondary antibody conjugated to DyLight® 633, which binds to hen immunoglobulins in immunological assays. DyLight® is a trademark of Thermo Fisher Scientific, Inc. and its subsidiaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.