

**German Research Products - GRP GmbH** 

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

## **Product Datasheet**

## Donkey anti-Rat IgG (H&L), DyLight® 550 conjugated GRP12812

Species/Host Donkey

Immunogen Purified rat IgG

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C. Product is stable for 4 weeks

at 2-8°C after rehydration. For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

**Note** For research use only.

**Clonality** Polyclonal

**Purity** Affinity purified donkey IgG

**Dilution Range** 1 : 20-1 : 2000 for most applications

**Application Notes** 

Additional Information: Based on immunoelectrophoresis, this antibody reacts with: heavy ( $\gamma$ ) chains on rat IgG light chains on all rat immunoglobulins.No reactivity is observed to: non-immunoglobulin rat serum proteins. Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. Background: Donkey anti-Rat IgG (H&L) - Affinity Pure, DyLight® 550 Conjugate is a secondary antibody conjugated to DyLight® 550, which binds to all rat IgG (H&L) in immunological assays. DyLight® 550 has Amax = 562 nm, Emax = 576 nm. Antibodies are are affinity purified using solid phase rat IgG (H&L).DyLight®

is a registered trade mark of Thermofisher Inc., and its subsidaries.

Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.