

German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

Donkey anti-Goat IgG (H&L), DyLight® 594 conjugated GRP12818

Species/Host Donkey

Immunogen Purified goat IgG, whole molecule

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C. Product is stable for 4 weeks

at 2-8°C after rehydration. For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

Note For research use only.

Clonality Polyclonal

Purity Affinity purified donkey IgG

Dilution Range 1 : 20-1 : 2000 for most applications

Application Notes Additional Information: Based on immunoelectrophoresis, this antibody reacts

with: heavy chains on goat IgG light chains on all goat immunoglobulins.No reactivity is observed to: non-immunoglobulin goat serum proteins.BSA and milk have to be replaced by other blocking reagents, like doneky serum or

commercial formulations which are free from bovine IgG. Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA,

Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. Background: Donkey anti-Goat IgG (H&L) , DyLight® 594 Conjugate is a secondary antibody conjugated to DyLight® 594, which binds to all goat IgGs in

immunological assays. DyLight® 550 has Amax = 593 nm, Emax = 618 nm. Antibodies are are affinity purified using solid phase goat IgG.DyLight® is a registered trade mark of Thermofisher Inc., and its subsidaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room

temperature to dissolve. Prepare fresh working dilutions daily.