

German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

Goat anti-Human kappa light chain, DyLight® 594 conjugated GRP12827

Species/Host Goat

Immunogen Purified human kappa light chain

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C. Product is stable for 4 weeks

at 2-8°C after rehydration. For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 0.5 mg of antibody in 0.55 ml of sterile water add 0.55 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to

mix well but without foaming.

Note For research use only.

Clonality Polyclonal

Purity Affinity purified goat IgG

Dilution Range 1 : 20-1 : 2000 for most applications

Application Notes Additional Information: Based on immunoelectrophoresis, this antibody reacts

with: kappa light chains on human immunoglobulins. No reactivity is observed to: non-immunoglobulin human serum proteins heavy chains on human immunoglobulins lambda light chains on human immunoglobulins. Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. Background: Goat anti-human kappa light chain , DyLight® 594 Conjugate is a secondary antibody conjugated to DyLight® 594, which binds to all human kappa light chain in immunological assays. DyLight® 550 has Amax = 593 nm, Emax = 618 nm. Antibodies are are affinity purified using solid phase human kappa light chain. DyLight® is a registered trade mark of Thermofisher Inc., and its subsidaries. Reconstitution: For reconstitution add 0.55 ml of sterile water.

Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.