

**German Research Products - GRP GmbH** 

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

## **Product Datasheet**

## Donkey anti-Goat IgG (H&L), DyLight® 633 conjugated GRP12852

Species/Host Donkey

**Immunogen** Purified goat IgG, whole molecule

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C. Product is stable for 4 weeks

at 2-8°C after rehydration. For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

**Note** For research use only.

**Clonality** Polyclonal

**Purity** Affinity purified donkey IgG

**Dilution Range** 1 : 20-1 : 2000 for most applications

**Application Notes** 

Additional Information: Based on immunoelectrophoresis, this antibody reacts with: chains on goat IgG light chains on all goat immunoglobulins. No reactivity is observed to: non-immunoglobulin goat serum proteins. BSA and milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free from bovine IgG. Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. Background: Donkey anti-Goat IgG (H&L) - DyLight® 633 Conjugated is a secondary antibody conjugated to DyLight® 633, which binds to goat IgG (H&L) in immunological assays. DyLight® 633 has Amax = 638 nm, Emax = 658 nm. Antibodies are are affinity purified using solid phase goat IgG. DyLight® is a registered trade mark of Thermofisher Inc., and its subsidaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve.

Prepare fresh working dilutions daily.