

**German Research Products - GRP GmbH** 

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

## **Product Datasheet**

## Rabbit anti-Hamster IgG (H&L), DyLight® 633 conjugated **GRP12854**

Species/Host Rabbit

Purified hamster IgG, whole molecule **Immunogen** 

Form/Appearance Lyophilized

Store lyophilized material at 2-8°C. Product is stable for 4 weeks Storage

at 2-8°C after rehydration. For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

Note For research use only.

Clonality Polyclonal

**Purity** Affinity purified rabbit IgG

**Dilution Range** 1:20-1:2000 for most applications

**Application Notes** Additional Information: Based on immunoelectrophoresis, this antibody reacts

with: (γ) chains on hamster IgG light chains on all hamster immunoglobulins.No

reactivity is observed to:non-immunoglobulin hamster serum proteins.

Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/lgG free. 0.05 % (w/v) sodium azide is added as preservative. Background: Rabbit anti-hamster IgG (H&L) - DyLight® 633 Conjugated is a secondary antibody conjugated to DyLight® 633, which binds to hamster IgG (H&L) in immunological assays. DyLight® 633 has Amax = 638 nm, Emax = 658 nm. Antibodies are are affinity purified using solid phase hamster IgG.DyLight® is a registered trade mark of Thermofisher Inc., and its

subsidaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working

dilutions daily.