

Product Datasheet

Goat anti-Human IgA (alpha chain), DyLight® 650 conjugated, min. cross-reactivity to bovine, mouse, rabbit serum GRP12882

| | |
|--------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Species/Host | Goat |
| Immunogen | Purified human IgA (alpha chain), whole molecule |
| Form/Appearance | Lyophilized |
| Storage | Store lyophilized material at 2-8°C. Product is stable for 4 weeks at 2-8°C after rehydration. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming. |
| Note | For research use only. |
| Clonality | Polyclonal |
| Purity | Affinity purified goat IgG |
| Dilution Range | 1 : 20-1 : 2000 for most applications |
| Application Notes | <p>Additional Information: Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. Based on immunoelectrophoresis, this antibody reacts with: heavy (α) chains on human IgA (α chain) No reactivity is observed to: non-immunoglobulin human serum proteins light chains on all human immunoglobulins serum proteins from bovine, mouse or rabbit</p> <p>Background: Goat anti-human IgA (α chain) - DyLight®650 Conjugated, min. cross-reactivity to bovine, mouse, rabbit serum is a secondary antibody conjugated to DyLight® 650, which binds to human IgA (α chain) in immunological assays. DyLight® 650 has Amax = 652 nm, Emax = 672 nm. Antibodies are affinity purified using solid phase human IgA (α chain). DyLight® is a registered trade mark of Thermofisher Inc., and its subsidiaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.</p> |