

German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

Goat anti-Human IgG (H&L), DyLight® 650 conjugated, min. cross-reactivity to bovine, mouse, rabbit serum GRP12886

Species/Host Goat

Immunogen Purified Human IgG, whole molecule

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C. Product is stable for 4 weeks

at 2-8°C after rehydration. For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

Note For research use only.

Clonality Polyclonal

Purity Affinity purified goat IgG

Dilution Range 1 : 20-1 : 2000 for most applications

Application Notes Ad

Additional Information: Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative.Based on immunoelectrophoresis, this antibody reacts with: heavy (γ) chains on human IgG. Light chains on all human immunoglobulinsNo reactivity is observed to: non-immunoglobulin human serum proteins serum proteins from bovine, mouse or rabbit Background: Goat anti-human IgG (H&L) - DyLight®650 Conjugated, min. cross-reactivity to bovine, mouse, rabbit serum is a secondary antibody conjugated to DyLight®650, which binds to human IgG (H&L) in immunological assays.DyLight®650 has Amax = 652 nm, Emax = 672 nm. Antibodies are are affinity purified using solid phase human IgG (H&L).DyLight® is a registered trade mark of

Thermofisher Inc., and its subsidaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve.

Prepare fresh working dilutions daily.