

## Product Datasheet

### Goat anti-Human kappa light chain, DyLight® 650 conjugated, min. cross-reactivity to Mouse serum GRP12890

<b>Species/Host</b>	Goat
<b>Immunogen</b>	Purified Human Kappa (κ) Chain
<b>Form/Appearance</b>	Lyophilized
<b>Storage</b>	Store lyophilized material at 2-8°C. Product is stable for 4 weeks at 2-8°C after rehydration. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 0.5 mg of antibody in 0.55 ml of sterile water add 0.55 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well but without foaming.
<b>Note</b>	For research use only.
<b>Clonality</b>	Polyclonal
<b>Purity</b>	Affinity purified goat IgG
<b>Dilution Range</b>	1 : 20-1 : 2000 for most applications
<b>Application Notes</b>	Additional Information: Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. Based on immunoelectrophoresis, this antibody reacts with: kappa (κ) light chains on human immunoglobulins. No reactivity is observed to: non-immunoglobulin human serum proteins heavy chains on human immunoglobulins lambda (λ) light chains on human immunoglobulins mouse serum proteins. Background: Goat anti-human kappa light chain - DyLight®650 Conjugated, min. cross-reactivity to mouse serum is a secondary antibody conjugated to DyLight® 650, which binds to human kappa light chain in immunological assays. DyLight® 650 has Amax = 652 nm, Emax = 672 nm. Antibodies are affinity purified using solid phase Human Kappa (κ) Chain. DyLight® is a registered trade mark of Thermofisher Inc., and its subsidiaries. Reconstitution: For reconstitution add 0.55 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.