

**German Research Products - GRP GmbH** 

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

## **Product Datasheet**

## Donkey anti-Sheep IgG (H&L), DyLight® 800 conjugated GRP12934

Species/Host Donkey

**Immunogen** Purified Sheep IgG, whole molecule

Form/Appearance Lyophilized

Storage Store lyophilized material at 2-8°C. Product is stable for 4 weeks

at 2-8°C after rehydration. For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

**Note** For research use only.

**Clonality** Polyclonal

**Purity** Affinity purified donkey IgG

**Dilution Range** 1 : 20-1 : 2000 for most applications

**Application Notes**Additional Information: Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v)

sodium azide is added as preservative.Based on immunoelectrophoresis, this antibody reacts with: heavy ( $\gamma$ ) chains on sheep IgG, light chains on all sheep immunoglobulinsNo reactivity is observed to: non-immunoglobulin sheep serum proteins.BSA and milk have to be replaced by other blocking reagents, like doneky serum or commercial formulations which are free from bovine IgG. Background: Donkey anti-Sheep IgG (H&L), DyLight® 800 Conjugated is a secondary antibody conjugated to DyLight® 800, which binds to Sheep IgG (H&L) in immunological assays.DyLight® 800 has Amax = 777 nm, Emax = 794 nm. Antibodies are purified using solid phase Sheep IgG (H&L)DyLight® is a registered trade mark of Thermofisher Inc., and its subsidaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room

temperature to dissolve. Prepare fresh working dilutions daily.