

**German Research Products - GRP GmbH** 

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

## **Product Datasheet**

## Donkey anti-Rabbit IgG (H&L), DyLight® 350 conjugated **GRP12941**

Species/Host Donkey

Purified Rabbit IgG, whole molecule **Immunogen** 

Form/Appearance Lyophilized

Store lyophilized material at 2-8°C. Product is stable for 4 weeks Storage

at 2-8°C after rehydration. For long time storage after

reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1.1 ml of sterile water add 1.1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard. Be sure to mix well

but without foaming.

Note For research use only.

Clonality Polyclonal

**Purity** Affinity purified donkey IgG

**Dilution Range** 1:20-1:2000 for most applications

**Application Notes** 

Additional Information: Conjugate is present in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05 % (w/v) sodium azide is added as preservative. Based on immunoelectrophoresis, this antibody reacts with: heavy  $(\gamma)$  chains on rabbit IgG, light chains on all rabbit immunoglobulinsNo reactivity is observed to: non-immunoglobulin rabbit serum proteins Background: Donkey anti-Rabbit IgG (H&L), DyLight®350 Conjugated is a secondary antibody conjugated to DyLight® 350, which binds to Rabbit IgG (H&L) in immunological assays.DyLight® 350 has Amax = 353 nm, Emax = 432 nm. Antibodies are purified using solid phase Rabbit IgG (H&L)DyLight® is a registered trade mark of Thermofisher Inc., and its subsidaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room

temperature to dissolve. Prepare fresh working dilutions daily.