

Product Datasheet

Goat anti-dog IgG (H&L), Ig fraction, Unconjugated GRP13004

| | |
|--------------------------|---|
| Species/Host | Goat |
| Reactivity | Dog |
| Immunogen | Purified Dog IgG, whole molecule. |
| Form/Appearance | Liquid |
| Storage | Store at 2-8 °C. |
| Note | For research use only. |
| Clonality | Polyclonal |
| Purity | Proprietary 2-step fractionation procedure |
| Dilution Range | The optimal working dilution should be determined by the investigator. |
| Application Notes | <p>Additional Information: Antibody purity is > 80% based on SDS-PAGE. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none">• heavy (γ) chains on dog IgG• light chains on all dog immunoglobulins <p>. Based on IEP, no reactivity is observed to:</p> <ul style="list-style-type: none">• non-immunoglobulin dog serum immunoglobulins <p>. Background: Goat anti-Dog IgG (H&L) Ig fraction is a secondary antibody which binds to Dog IgG (H&L) in immunological assays</p> |