

Product Datasheet

Goat anti-Donkey IgG (H&L), Biotin conjugated GRP13006

Species/Host	Goat
Reactivity	Donkey
Tested Applications	FC, IL
Immunogen	Purified Donkey IgG, whole molecule.
Form/Appearance	Liquid
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Dilution Range	1 : 20-1 : 2000 (Flow cyt), (IL)
Application Notes	<p>Additional Information: Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Donkey IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none">• heavy (γ) chains on donkey IgG• light chains on all donkey immunoglobulins <p>Based on IEP, no reactivity is observed to:</p> <ul style="list-style-type: none">• non-immunoglobulin donkey serum immunoglobulins <p>Background: Goat anti-Donkey IgG (H&L) is a secondary antibody, conjugated to biotin, which binds to Donkey IgG (H&L) in immunological assays.</p>