

Product Datasheet

Goat anti-Donkey IgG (H&L), DyLight® 633 conjugated GRP13007

Species/Host	Goat
Reactivity	Donkey
Tested Applications	FC, IL
Immunogen	Purified Donkey IgG, whole molecule.
Form/Appearance	Lyophilized
Storage	Store freeze-dried powder at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Dilution Range	1 : 20-1 : 2000 (Flow cyt), (IL)
Application Notes	<p>Additional Information: Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Donkey IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none">• heavy (γ) chains on donkey IgG• light chains on all donkey immunoglobulins <p>Based on IEP, no reactivity is observed to:</p> <ul style="list-style-type: none">• non-immunoglobulin donkey serum immunoglobulins <p>Background: Goat anti-Donkey IgG (H&L), DyLight® 633 conjugated is a secondary antibody conjugated to DyLight® 633, which binds to Donkey IgG (H&L) in immunological assays. DyLight® 633 has Amax = 638 nm, Emax = 658 nm. DyLight® is a registered trade mark of Thermofisher Inc., and its subsidiaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.</p>