

## Product Datasheet

### Rabbit anti-Goat IgG (H&L), DyLight® 350 conjugated GRP13010

Species/Host	Rabbit
Reactivity	Goat
Tested Applications	FC, IL
Immunogen	Purified Goat IgG, whole molecule.
Form/Appearance	Lyophilized
Storage	Store freeze-dried powder at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Dilution Range	1 : 20-1 : 2000 (Flow cyt), (IL)
Application Notes	<p>Additional Information: Antibody purity is &gt; 95% based on SDS-PAGE. Affinity purified using solid phase Goat IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: • heavy (γ) chains on goat IgG • light chains on all goat immunoglobulins. Based on IEP, no reactivity is observed to: • non-immunoglobulin goat serum immunoglobulins. Background: Rabbit anti-goat IgG (H&amp;L), DyLight®350 Conjugated is a secondary antibody conjugated to DyLight® 350, which binds to Goat IgG (H&amp;L) in immunological assays. DyLight® 350 has Amax = 353 nm, Emax = 432 nm. Antibodies are purified using solid phase Goat IgG (H&amp;L). DyLight® is a registered trade mark of Thermofisher Inc., and its subsidiaries. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.</p>