

Product Datasheet

Goat anti-Human IgG Fc, min. cross-reactivity to human IgA + IgM, bovine, horse, mouse or rabbit serum proteins GRP13039

Species/Host	Goat
Reactivity	Human
Immunogen	Purified Human IgG, Fc fragment
Form/Appearance	Liquid
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Uniprot ID	O75015
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Human IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: • heavy (γ) chains on human IgG. Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgA or IgM • serum proteins from bovine, horse, mouse, or rabbit.</p> <p>Background: Goat anti-Human IgG Fc, min. cross-reactivity to bovine, horse, mouse or rabbit serum proteins or mouse IgG1 is a secondary antibody which binds to human IgG Fc in immunological assays.</p>