German Research Products - GRP GmbH
In der Stockwiese 26
D-85410 Haag/Amper, Germany
Email: info@grp-ak.de
Phone: +49 (0)8167 6335

Product Datasheet

Goat anti-Human IgG Fc, TRITC conjugated, min. cross-reactivity to bovine, horse, mouse or rabbit serum proteins or mouse IgG1

GRP13040

Species/Host	Goat
Reactivity	Human
Tested Applications	FC, IL
Immunogen	Purified Human IgG, Fc fragment
Form/Appearance	Lyophilized
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Uniprot ID	O75015
Dilution Range	1 : 20-1 : 2000 (Flow cyt), (IL)
Application Notes	Additional Information: Antibody purity is > 95% based on SDS-PAGE.Affinity purified using solid phase Human IgG .Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative.Based on IEP, this antibody reacts with: • heavy (γ) chains on human IgG .Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgG F(ab)'2 fragment • mouse IgG1 • serum proteins from bovine, horse, mouse, or rabbit . Background: Goat anti-Human IgG Fc, TRITC conjugated, min. cross-reactivity to bovine, horse, mouse or rabbit serum proteins or mouse IgG1 is a secondary antibody conjugated to TRITC, which binds to human IgG Fc in immunological assays. Reconstitution: For reconstitution add 1.1 ml of sterile water.Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.