

Product Datasheet

Goat anti-Human IgG Fc, TRITC conjugated, min. cross-reactivity to human IgA + IgM, bovine, horse, mouse or rabbit serum proteins GRP13041

Species/Host	Goat
Reactivity	Human
Tested Applications	FC, IL
Immunogen	Purified Human IgG, Fc fragment
Form/Appearance	Lyophilized
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Uniprot ID	O75015
Dilution Range	1 : 20-1 : 2000 (Flow cyt), (IL)
Application Notes	<p>Additional Information: Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Human IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none">• heavy (γ) chains on human IgG• Based on IEP, no reactivity is observed to:• non-immunoglobulin human serum immunoglobulins• light chains on all human immunoglobulins• human IgA or IgM• serum proteins from bovine, mouse, or rabbit <p>Background: Goat anti-Human IgG Fc, TRITC conjugated, min. cross-reactivity to human IgA + IgM, bovine, horse, mouse or rabbit serum proteins is a secondary antibody conjugated to TRITC, which binds to human IgG Fc in immunological assays. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.</p>