

Product Datasheet

Goat anti-Human IgG Fc, F(ab)'2 Fragment, min. cross-reactivity to Human IgA/IgM, Bovine, Horse, Mouse, Rabbit Serum GRP13042

Species/Host	Goat
Reactivity	Goat
Immunogen	Purified Human IgG, Fc
Form/Appearance	Liquid
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: Antibody purity is > 90% based on SDS-PAGE Small amounts of intact IgG may be present. Affinity purified using solid phase Human IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none">• heavy (γ) chains on human IgG• Based on IEP, no reactivity is observed to:• non-immunoglobulin human serum immunoglobulins• light chains on all human immunoglobulins• human IgA or IgM• serum proteins from bovine, mouse, or rabbit <p>Background: Goat anti-Human IgG Fc, F(ab)'2 Fragment, min. cross-reactivity to Human IgA/IgM, Bovine, Horse, Mouse, Rabbit Serum is a secondary antibody, which binds to human IgG Fc, F(ab)'2 Fragment in immunological assays.</p>