

## Product Datasheet

### Rabbit anti-Human IgM ( $\mu$ chain), F(ab)'<sub>2</sub> Fragment GRP13055

Species/Host	Rabbit
Reactivity	Human
Immunogen	Llama IgG
Form/Appearance	Lyophilized
Storage	Store at 2 - 8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Dilution Range	The optimal working dilution should be determined by the investigator.

**Application Notes** Additional Information: Antibody purity is > 90 % based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Human IgM. F(ab)'<sub>2</sub> fragment was prepared by pepsin digestion. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. Based on IEP, this antibody reacts with: • heavy ( $\mu$ ) chains on Human IgM. Based on IEP, no reactivity is observed to: • light chains on all Human immunoglobulins • non-immunoglobulin Human serum proteins • Mouse IgG or serum proteins. Background: Rabbit anti-Human IgM ( $\mu$  chain), F(ab)'<sub>2</sub> Fragment is a secondary antibody, which binds to human IgM ( $\mu$  chain), F(ab)'<sub>2</sub> Fragment in immunological assays. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.