

Product Datasheet

Goat anti-Human IgM (μ chain), F(ab)'2 Fragment, min. cross-reactivity to IgG from human or mouse GRP13056

| | |
|-----------------|--|
| Species/Host | Goat |
| Reactivity | Goat |
| Immunogen | Purified Human IgM, μ chain |
| Form/Appearance | Liquid |
| Storage | Store at 2-8 °C. |
| Note | For research use only. |
| Clonality | Polyclonal |
| Purity | Affinity purified |
| Uniprot ID | AAC37537.1 |
| Dilution Range | The optimal working dilution should be determined by the investigator. |

Application Notes Additional Information: Antibody purity is > 90% based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Human IgM. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: • heavy (γ) chains on human IgM. Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum proteins • light chains on all human immunoglobulins • IgG from human or mouse. Background: Goat anti-Human IgM (μ chain), F(ab)'2 Fragment, min. cross-reactivity to IgG from human or mouse is a secondary antibody, which binds to human IgM (μ chain) in immunological assays.