

## **German Research Products - GRP GmbH**

In der Stockwiese 26 D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

## **Product Datasheet**

## Goat anti-Mouse IgG (H&L), DyLight® 488 conjugated, min. cross-reactivity to bovine, horse, human, pig or rabbit serum proteins GRP13062

**Species/Host** Goat

Reactivity Mouse

Tested Applications FC, IL

**Immunogen** Purified Mouse IgG, whole molecule

Form/Appearance Lyophilized

**Storage** Store freeze-dried powder at 2-8 °C.

**Note** For research use only.

**Clonality** Polyclonal

**Purity** Affinity purified

Uniprot ID AAA51107

**Dilution Range** 1 : 20-1 : 2000 (Flow cyt), (IL)

**Application Notes** 

Additional Information: Antibody purity is > 95% based on SDS-PAGE.Affinity purified using solid phase Mouse IgG .Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative.Based on IEP, this antibody reacts with: • heavy ( $\gamma$ ) chains on mouse IgG • light chains on all mouse

immunoglobulins .Based on IEP, no reactivity is observed to: •

non-immunoglobulins .based of IEF, no reactivity is observed to. on non-immunoglobulin mouse serum immunoglobulins • serum proteins from bovine, horse, human rabbit or swine . Background: Goat anti-Mouse IgG (H&L), DyLight® 488 conjugated, min. cross-reactivity to bovine, horse, human, pig or rabbit serum proteins is a secondary antibody conjugated to DyLight® 488, which binds to mouse IgG (H&L) in immunological assays.DyLight® 488 has Amax = 493 nm, Emax = 518 nm.DyLight® is a registered trade mark of Thermofisher Inc., and its subsidaries. Reconstitution: For reconstitution add 1.1 ml of sterile water.Let it stand 30 minutes at room temperature to dissolve.

Prepare fresh working dilutions daily.