

## Product Datasheet

### Goat anti-Mouse IgG (H&L), F(ab)'2 Fragment, min. cross-reactivity to bovine, horse, human, pig or rabbit serum protein GRP13088

Species/Host	Goat
Reactivity	Mouse
Immunogen	Purified Mouse IgG, whole molecule
Form/Appearance	Liquid
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Uniprot ID	AAA51107
Dilution Range	The optimal working dilution should be determined by the investigator.

**Application Notes** Additional Information: Antibody purity is > 90% based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Mouse IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: • heavy (γ) chains on mouse IgG • light chains on all mouse immunoglobulins. Based on IEP, no reactivity is observed to: • non-immunoglobulin mouse serum immunoglobulins • serum proteins from bovine, horse, human rabbit or swine. Background: Goat anti-Mouse IgG (H&L), F(ab)'2 Fragment, min. cross-reactivity to bovine, horse, human, pig or rabbit serum protein is a secondary antibody which binds to mouse IgG (H&L), F(ab)'2 Fragment in immunological assays.